

Pediatric Dosing Chart

TYLENOL (ACETAMINOPHEN)

WEIGHT	ELIXIR 160MG/5ML	CHEWABLE TABS 80MG/TABLET	JR. STRENGTH CAPS 160MG/CAPSULE
6-11 LBS	1/4 TSP (1.25ML)		
12-17 LBS	1/2 TSP (2.5ML)		
18-23 LBS	3/4TSP (3.5ML)		
24-35 LBS	1 TSP (5ML)	2 TABS	
36-47 LBS	1 1/2 TSP (7.5ML)	3 TABS	
48-59 LBS	2 TSP	4 TABS	2 CAPS
60-71 LBS	2 1/2 TSP	5 TABS	2 1/2 CAPS
72-95 LBS	3 TSP	6 TABS	3 CAPS

May give one dose every four hours.

BENADRYL (DIPHENHYDRAMINE)

WEIGHT	LIQUID 12.5MG/TSP	CHEWABLES 12.5MG	TABS/CAPSULES 25MG
22-32 LBS	1 TSP (5ML)	1 CHEWABLE	
33-42 LBS	1 1/2 TSP (7.5ML)	1 1/2 CHEWABLE	
43-53 LBS	2 TSP (10ML)	2 CHEWABLE	1 TAB/CAPSULE

May give one dose every six hours.

Ibuprofen (Motrin, Advil) should NOT be used under 6 months of age. After 6 months of age, the dosage is determined by weight.

MOTRIN/ADVIL (IBUPROFEN)

WEIGHT	INFANT DROPS 50MG/1.25ML	CHILDREN'S LIQUID 100MG/TSP	JR. CHEWABLE 100MG
12-17 LBS	1.25 ML		
18-23 LBS	1.875 ML		
24-35 LBS	2.5 ML	1 TSP (5ML)	1 TAB
36-47 LBS		1 1/2 TSP (7.5ML)	1 1/2 TABS
48-59 LBS		2 TSP (10ML)	2 TABS

May give a dose every six to eight hours.

**Never give your child a medication that they've had a previous reaction to.*